

## § 520.1205

milligrams of kanamycin as the sulfate, 0.033 milligram of aminopentamide hydrogen sulfate, 25 milligrams of pectin, 250 milligrams of bismuth subcarbonate, and 500 milligrams of activated attapulgite.

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) It is administered orally to dogs for the symptomatic relief of acute bacterial diarrhea caused by kanamycin-susceptible organisms.

(2) The drug is recommended for use at the rate of one teaspoonful (5 milliliters) of suspension per 20 pounds of body weight every 8 hours. Animals weighing under 10 pounds should be given one-half the above amount every 8 hours. The initial dose should be twice the amount of a single dose. Maximum dosage should not exceed three times the recommended dose.

(3) For use only by or on the order of a licensed veterinarian.

[40 FR 13838, Mar. 27, 1975, as amended at 53 FR 27851, July 25, 1988; 56 FR 8710, Mar. 1, 1991; 64 FR 403, Jan. 5, 1999]

## § 520.1205 Kanamycin sulfate, pectin, bismuth subcarbonate, activated attapulgite tablets.

(a) *Specifications.* Each tablet contains 100 milligrams of kanamycin (as the sulfate), 25 milligrams of pectin, 250 milligrams of bismuth subcarbonate, and 500 milligrams of activated attapulgite.

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount.* One tablet per 44 kilograms (20 pounds) of body weight every 8 hours. Maximum dose 3 tablets every 8 hours. For animals under 22 kilograms (10 pounds)  $\frac{1}{2}$  tablet every 8 hours. The initial loading dose should be twice the amount of a single dose.

(2) *Indications for use.* For the treatment of bacterial enteritis caused by organisms susceptible to kanamycin and the symptomatic relief of associated diarrhea in dogs.

(3) *Limitations.* Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[56 FR 8710, Mar. 1, 1991]

## 21 CFR Ch. I (4–1–05 Edition)

### § 520.1242 Levamisole hydrochloride oral dosage forms.

#### § 520.1242a Levamisole powder for oral solution.

(a) *Specifications.* Each package of powder contains 9.075, 11.7, 18.15, 46.8, 362.7, or 544.5 grams (g) levamisole hydrochloride.

(b) *Sponsors.* See sponsors in § 510.600(c) for use as follows:

(1) No. 000061 for use of 46.8- and 544.5-g packages as in paragraph (e)(1)(i), (e)(1)(ii)(B), and (e)(1)(iii) of this section; for 11.7-, 46.8-, and 544.5-g packages as in paragraph (e)(2)(i), (e)(2)(ii)(B), and (e)(2)(iii) of this section; and for an 18.15-g package as in paragraph (e)(3) of this section.

(2) No. 053501 for use of a 46.8-g package as in paragraph (e)(1)(i), (e)(1)(ii)(A), and (e)(1)(iii) of this section; for 11.7- and 46.8-g packages as in paragraph (e)(2)(i), (e)(2)(ii)(A), and (e)(2)(iii) of this section; and for 9.075- and 18.15-g packages as in paragraph (e)(3) of this section.

(3) No. 057561 for use of 46.8- and 544.5-g packages as in paragraphs (e)(1)(i), (e)(1)(ii)(A), and (e)(1)(iii) and (e)(2)(i), (e)(2)(ii)(A), and (e)(2)(iii) of this section.

(4) No. 059130 for use of 46.8-, 362.7-, and 544.5-g packages as in paragraphs (e)(1)(i), (e)(1)(ii)(B), (e)(1)(iii), (e)(2)(i), (e)(2)(ii)(B), and (e)(2)(iii) of this section; and for use of an 18.15-g package as in paragraph (e)(3) of this section.

(c) *Related tolerances.* See § 556.350 of this chapter.

(d) *Special considerations.* See § 500.25 of this chapter.

(e) *Conditions of use.* It is used as an anthelmintic as follows:

(1) *Cattle*—(i) *Amount.* 8 milligrams per kilogram (mg/kg) body weight as a drench.

(ii) *Indications for use*—(A) Effective against the following nematode infections: Stomach worms (*Haemonchus*, *Trichostrongylus*, *Ostertagia*); intestinal worms (*Trichostrongylus*, *Cooperia*, *Nematodirus*, *Bunostomum*, *Oesophagostomum*); and lungworms (*Dictyocaulus*).

(B) Effective against the following adult nematode infections: Stomach worms (*Haemonchus placei*, *Ostertagia*